

Comparative study of cataract in hypertensive patients and non-hypertensive patients

Reshma Mehta^{1,*}, Manjunath Patil², Shubashree Page³

^{1,2}Assistant Professor, ³JR, Dept. of Ophthalmology, Ashwini Rural Medical College, Solapur, Maharashtra

***Corresponding Author:**
Email: dradmehta@gmail.com

Abstract

Introduction: The purpose behind conducting this study was to determine if hypertension increases the risk of cataract, to find out common morphological types of cataract in hypertensive patients, to determine relation between grade of hypertension and grade of nuclear sclerosis and to compare prevalence rates of cataract in hypertensive patients on anti-hypertensive drugs and those not on anti-hypertensive drugs.

Materials & Methods: Comparative study of cataract in hypertensive patients and non-hypertensive patients was carried out in Department of ophthalmology at Ashwini Rural Medical College, Kumbhari, Solapur with 300 eligible participants.

Results: This was a comparative study with 156 hypertensive and 144 non-hypertensive patients. Prevalence of cataract was found to be 79.53%. Prevalence of cataract was significantly higher in eyes of hypertensive patients as compared to eyes of non-hypertensive patients; (87.83% and 70.8% respectively, with p-value<0.0001). Significant association was found between high systolic BP and cataract. ($\chi^2=23.82$, df= 1, p<0.0001). Nuclear cataract was the most common morphological type in eyes of hypertensive patients (50.4%). However, posterior sub capsular cataract was more common in eyes of hypertensive patients as in eyes of non-hypertensive patients (28.6% and 23.3%). No significant relation was found between grade of hypertension and grade of nuclear sclerosis ($\chi^2 = 4.33$, df= 3, p=0.23). No significant difference between them was found in our study. (Z=0.10, p=0.92).

Conclusions: Risk of cataract is higher in hypertensive patients as compared to non-hypertensive patients. Relation between grade of hypertension and grade of nuclear sclerosis and role of anti-hypertensive drugs must be further explored.

Keywords: Cataract, Hypertension, Non-hypertensive patients, Risk factors, Prevalence

| Access this article online | |
|---|---|
| Quick Response Code: | Website: www.innovativepublication.com |
|  | DOI: 10.5958/2395-1451.2016.00034.2 |

Introduction

Blindness is a public health problem worldwide. According to WHO global data for visual impairments 2010, prevalence of blindness was 20.5%¹. The prevalence rate is significantly higher in India when compared to Europe (7%) and America (8%). Almost all of the blindness (80%) is preventable^{1,2}. Cataract is defined as any opacity in lens or capsule whether congenital or acquired. Cataract is the leading cause of blindness worldwide³. According to National Survey on blindness 2006-07, cataract is responsible for about 62.6% of all cases of blindness in India⁴. According to WHO cataract is responsible for nearly 50% of cases of blindness worldwide^{1,5}. Nearly 12 to 15 million individuals are thought to be blind by cataract worldwide. Cataract occurs a decade earlier in India than the Western world³. According to Rapid Assessment of Avoidable Blindness-INDIA8 Report 2006-07, (RAABI) 77.5 percent of blindness (vision <6/60) is caused by cataract⁵. The prevalence

of blindness was observed to be 1.34 times higher in females compared to males.^{5,6,7,8,9} It is also found that prevalence of blindness above 70 years to be 16 times higher than those aged 50-54 years¹⁰. Currently, the only treatment available for cataract is surgical extraction of the lens¹¹. Hence it is important to study the possible risk factors for cataract. A better and detailed understanding of etio-pathogenesis of cataract can help us take appropriate measures to delay the onset of senile cataract. Hypertension is diagnosed by average awake systolic blood pressure ≥ 140 mmHg and/or diastolic blood pressure ≥ 90 mmHg according to American Heart Association¹². According to a community based survey carried out by ICMR during 2007-08 prevalence of hypertension in India was 17 to 21% with marginal rural urban differences. Overall pattern of prevalence was found to be increasing with all age groups⁴. Hence it is important to study the effects of hypertension on the lens due to increasing morbidity of hypertension. Hypertension is known to be associated with ocular conditions like hypertensive retinopathy, ocular hypertension, etc.¹³. Various studies have been carried out in the past to study the relation between hypertension and cataract; however the need for further research has been put forth by most of them. Some of the previous epidemiological studies¹⁴⁻²⁵ indicate that hypertension plays an important role in development of cataract while some others do not.^{11,26,27,28}

Patho-physiologic mechanisms of association of hypertension with cataract are however unclear and role

of oxidative stress, IL6, TNF α and elevated levels of C-reactive protein have been suggested²⁹. Some of the past studies also state that associations may be due to confounding factors like smoking or diabetes mellitus²⁶. Some epidemiological studies state that there is increased risk of cataract with high systolic but not diastolic blood pressure²⁶. Some epidemiological studies have concluded that hypertension acts as a significant risk factor for Posterior sub-capsular cataract (PSC)¹⁹. Association between nuclear and cortical cataract and hypertension gave varied results in different epidemiological studies. Some studies suggest that hypertension is significantly associated with nuclear cataract²⁵, while some studies have concluded that hypertension is not significantly associated with nuclear cataract¹⁹. Results for association of cortical cataract with hypertension were found to be inconclusive by some studies¹⁹. Some of the previous studies have also concluded that people with severe hypertension had a greater risk of cataract¹⁹. The results of various studies of role of anti-hypertensive drugs and occurrence of cataract are conflicting. Some studies suggest that use of anti-hypertensive drugs can paradoxically promote cataract formation^{30,31,32,33}, while some others conclude that anti-hypertensive drugs have a protective effect on the lens³². Some other studies on the other hand suggest that use of anti-hypertensive drugs is not associated with any type of cataract³⁰.

The purpose behind conducting this study was to determine an independent relation between hypertension and cataract by eliminating major confounding factors like Diabetes Mellitus, smoking and alcoholism. Not many studies with respect to this subject have been carried out in the state of Maharashtra.

Materials and Methods

This comparative study was carried out according to the ethical principles for medical research involving human subjects defined by the 'Declaration of Helsinki'. An approval from the Institutional Ethics Committee (IEC) was obtained before starting the study. Study was done in department of Ophthalmology at Ashwini Rural Medical College, Kumbhari, Solapur a tertiary care hospital in Solapur district, during two months duration. Patients who fulfilled our inclusion criteria were included in the study, accordingly a total 300 patients were included.

Inclusion criteria:

1. Patients with age ≥ 50 years visiting OPD of ophthalmology or general medicine of our institute
2. Patients who are willing to participate in the study.

Exclusion criteria:

1. Patients who do not give consent for this study.
2. Patients with age < 50 years.
3. Diabetic patients (Random blood sugar > 200 mg/dl)³⁴ as well as known cases of

Diabetes Mellitus.

4. Patients who are smokers and/or consume alcohol.

This was a comparative study consisting of 300 patients. One group was classified as 'hypertensive' and the other group being 'non-hypertensive'.

Systolic and diastolic blood pressure of all eligible patients was measured (after allowing them to rest for at least five minutes before measuring blood pressure) using a mercury sphygmomanometer in right arm sitting position using a cuff of appropriate size. The blood pressure recorded was graded as follows.^{34,35}

| Blood Pressure classification | Systolic mmHg | Diastolic mmHg |
|-------------------------------|---------------|----------------|
| Normal | <120 | And <80 |
| Pre-hypertension | 120-139 | Or 80-89 |
| Stage 1 | 140-159 | Or 90-99 |
| Stage 2 | >160 | Or >100 |
| Isolated systolic | >140 | And <90 |

Statistical analysis

Descriptive statistics such as mean, SD and percentage were used. Comparison between groups was done by using z-test for proportion. Chi-square test was used to find out association between categorical variables. A p-value less than 0.05 were considered as significant. Data was analyzed using MS-excel.

Results

At the end of study duration, we got 300 participants for our study who were examined and interviewed for all the parameters in our questionnaire.

Table 1: Distribution of hypertensive and non-hypertensive cases

| | No. of cases | Percentage |
|------------------|--------------|------------|
| Hypertensive | 156 | 52 |
| Non-hypertensive | 144 | 48 |
| Total | 300 | 100 |

Out of 300 patients, 156 (52%) were hypertensive and 144 (48%) were non-hypertensive.

Table 2: Prevalence of cataract in 'eyes' of study population

| | No. of cases | Percentage |
|------------|--------------|------------|
| Cataract | 408 | 79.53 |
| Clear lens | 105 | 20.47 |
| Total | 513 | 100 |

Total 600 eyes of 300 patients were examined. We excluded 'eyes' previously operated for cataract and corneal opacities. Hence, we examined total 513 'eligible eyes'. Out of these, cataract was detected in 408 eyes. Hence prevalence of cataract was 79.53%.

Table 3: Comparison of prevalence of cataract in eyes of hypertensive and non-hypertensive patients

| Cataract | Eyes of Hypertensive Patients (%) | Eyes of Non-hypertensive Patients (%) | Total |
|----------|-----------------------------------|---------------------------------------|-------|
| Present | 231 (87.83) | 177 (70.8) | 408 |
| Absent | 32 (12.17) | 73 (29.2) | 105 |
| Total | 263 | 250 | 513 |

Z=4.85, p<0.0001

Out of 513 eyes, 263 eyes were of hypertensive patients and 250 eyes were of non-hypertensive patients. Cataract was seen in 231 eyes of hypertensive patients and 177 eyes of non-hypertensive patients. Hence, Prevalence of cataract in eyes of hypertensive patients was 87.83% (231/263) Prevalence of cataract in eyes of non-hypertensive patients was 70.8% (177/250).

From the above table, prevalence of cataract was observed more in eyes of hypertensive patients (87.83%) than eyes of non-hypertensive patients (70.8%) and also found that, there is highly statistical significant difference between prevalence of cataract in eyes of hypertensive and non- hypertensive patients (p<0.0001). Hence our research hypothesis was accepted.

Table 4: Morphological types of cataract in eyes of hypertensive and non-hypertensive patients

| Type of morphology | Eyes of HTN patients (%) | Eyes of Non-HTN patients (%) | Total |
|-----------------------|--------------------------|------------------------------|-------|
| Anterior capsular | 3 (0.9) | 1 (0.4) | 4 |
| Brown cataract | 6 (1.8) | 7 (2.8) | 13 |
| Cortical | 31 (9.1) | 27 (10.7) | 58 |
| Nuclear | 171 (50.4) | 127 (50.19) | 298 |
| PSC | 97 (28.6) | 59 (23.3) | 156 |
| Total mature cataract | 22 (6.5) | 24 (9.5) | 46 |
| Hyper mature cataract | 0 (0) | 1 (0.4) | 1 |
| Total | 330 (100) | 246 (100) | 576 |

Based on the morphology of cataract detected on slit lamp examination, we determined the most common morphological type of cataract in hypertensive patients. In cases of mixed cataracts, each of the morphological types was considered separately.

It is observed that, Nuclear cataract was found to be most common morphological type of cataract in eyes of hypertensive patients (171 eyes; either purely nuclear or in combination with other morphological types). This was followed by posterior sub capsular (PSC; 28.6%) and cortical cataract (9.1%).

Similar pattern was seen in non-hypertensive patients; however, prevalence of posterior sub- capsular cataract was higher in hypertensive patients than non-hypertensive patients. (28.6% against 23.3%).

Discussion

Cataract and hypertension: From the above results, we have found that prevalence of cataract is significantly higher in hypertensive patients (87.83%) as compared to non-hypertensive patients (70.8%). This difference in prevalence is independent of confounding factors such as diabetes mellitus, smoking and alcoholism. Hence, our first research hypothesis i.e. 'There is significant difference in the prevalence rates of cataract in hypertensive patients and non-hypertensive patients' is

supported. Hence, risk of cataract is more in hypertensive people than non-hypertensive people. M Sharma, et. al²⁷ also found a higher prevalence of cataract in hypertensives (74%) as compared to non-hypertensives (69.8%). However, this difference was not found to be statistically significant. Similar to our study, many of the previous studies found hypertension as a significant risk factor for development of cataract¹⁴⁻²⁵.

Consistent with results by Schaumberg, et. al²⁶, Xiaoning Yu, et.al¹⁹, we have found high systolic blood pressure as a risk factor for cataract.

Some studies^{11,26,27,28} however did not find hypertension as a risk factor with cataract.

Patho-physiological mechanisms of role of hypertension in development of cataract is not clear. A study by Bautista LE, et. al²⁹; has found that elevated plasma levels of IL-6 and TNF-a appeared in individuals with high blood pressure (Systolic BP>=140mmHg or diastolic BP>=90mmHg). A study by Virgolic B, et. al³⁶ revealed that senile cataract is a systemic disease with an inflammatory component. Hence, role of hypertension in development of cataract can possibly be explained by raised inflammatory markers in hypertension.

A study by J Kaur, et. al³⁷ have found that oxidative stress plays an important role in onset and progression

of cataract. In a study by Briones AM, et. al.³⁸ role of oxidative stress in hypertension was found. Hence, role of hypertension in development of cataract can be attributed to oxidative stress.

In a study by Flipovsky et. al.³⁹ it was found in their studies that cortisol levels are significantly higher in men with high systolic blood pressure than diastolic blood pressure. This can possibly explain association of high systolic blood pressure with cataract; as corticosteroids are known to cause cataract³.

Lee SM, et. al.⁴⁰ have found in their study that systemic hypertension induces change in the protein conformational structures of the lens capsule, then cause altered membrane transport and permeability of ions and result in exacerbation of cataract formation. This could be one of the mechanisms by which hypertension plays a role in cataract formation.

A study by Smith PA, et. al.⁴¹ has revealed that sympathetic hyperactivity play an integral role in the development of hypertension and its complications. Similar results were found by Masuo K, et. al.⁴² who conducted studies on heightened sympathetic nervous system activity contribute to elevation of blood pressure.

Churchill GC, et. al.⁴³ have reported presence of α 1A adrenergic receptors in the lens. It is not known if these play a role in cataract formation, however the role of these can be explored in the future.

Common morphology of cataract: We found nuclear cataract most common morphological type in our study. This was seen in both hypertensive as well as non-hypertensive patients. This finding of our study is consistent with study by Tsai, et. al.²¹.

The prevalence rates of nuclear cataracts were however nearly equal in hypertensive and non-hypertensive patients. (50.4% and 50.19% respectively).

Posterior sub-capsular was the next common type. The prevalence of posterior sub capsular cataract was higher in hypertensives (28.6%) as compared to non-hypertensives (23.3%). Tsai, et. al.²¹ however found prevalence of cortical cataract higher than that of posterior sub capsular cataract in their study (12.9% and 9.2% respectively). This can probably be so as we had excluded diabetic patients from our study. A Study by Neil Rowe, et.al⁴⁴ had found diabetes as a risk factor for cortical cataract.

Higher prevalence of posterior sub capsular cataract in hypertensive patients can possibly be due to higher level of corticosteroids in hypertensive patients as compared to non-hypertensive patients. Harold Skalka et. al⁴⁵ have found posterior sub capsular cataract to be significantly associated with corticosteroids.

Conclusion

From our study, we have concluded that prevalence of cataract is significantly higher in hypertensive patients as compared to non-hypertensive patients. There is significant association of high systolic

blood pressure with cataract. Hence, high systolic blood pressure acts as a significant risk factor for cataract. Also, there is increased risk of cataract in hypertensive patients as compared to non-hypertensive patients.

Nuclear cataract is the most common morphological type seen in hypertensive patients; however there was a difference in prevalence of posterior sub capsular cataract in eyes of hypertensive and non-hypertensive patients (28.6% and 23.3%). The relation between grade of hypertension and grade of nuclear hardness (nuclear sclerosis) was not found to be significant in our study. Further research will be needed on this topic after matching for factors like age.

References

1. Rensikoff S, Pascolini D, et.al, Kocurl, et.al(2004) Global data on visual impairment in year 2002. Bull world health organ 82:844-851;doi:10.1590/S0042-9686200400110009.
2. Limburg H, World Health Organisation for rapid assessment of cataract surgical services. Prevention of blindness and deafness, Geneva, WHO, Switzerland 2001;1-604.
3. Parson's disease of eye Ramanjit Sihota, Radhika Tandon 22nd edition 262-227, 98.
4. K. Park; Park's textbook of preventive and social medicine 23rd edition 374,402.
5. Directorate of General Health Services, National Program for Control of Blindness, Rapid assessment of Avoidable Blindness-INDIA report DGHS,MOHFW, GOI, New Delhi 2006-2007;1-56.
6. Muralikrishnan R, Krishna PR, Thulasiraj RD, Bachani B, Guptha S et.al., Blindness estimation s, projections and service delivery NCMH background papers- Burden of disease in India 2006;305-17.
7. Murthy GVS, Gupta KS, John N, Vashist P Current status of Cataract blindness and Vision 2020: the right to sight initiative in India. Indian J Ophthalmol 2008;56:489-94.
8. National Survey on Blindness and visual outcomes after cataract surgery (2001-2002) Report. National programme for control of Blindness, MOHFW,G O I 2002.
9. Neena J, Rachel J, Praveen V, Murthy GVS Rapid assessment of avoidable blindness in India PLoS One 2008;3:e2867,www.plosone.org.
10. WHO global data on visual impairments 2010.
11. Survey of ophthalmology volume 39. Number 4. January-February 1995; Public health and the eye; Epidemiology of risk factors for age-related cataract; Sheila K. West, et.al.
12. Roger VL, et.al; Heart disease and stroke statistics— 2012 update: a report from the American Heart Association. Circulation. 2012 Jan 3.125(1):e2-e220.
13. Wang S, Xu L, Jonas JB, et al. major eye diseases and risk factors associated with systemic hypertension in an adult Chinese population; the Beijing eye study. Ophthalmology 2009;116:2373-80.
14. Diabetes, Hypertension and central obesity as risk factors as cataract risk factors in a black population, The Barbados Eye Study, Leske MC, et.al; Ophthalmology. 1999 Jan;106(1):35-41.
15. Metabolic syndrome components and Age related cataract: The Singapore Malay Eye Study; Sabanayagam,et.al; Invest Ophthalmol Vis Sci

- 01/2011;52:2397-2404.
16. Association between metabolic syndrome components and age related cataract; S. Park, et.al; Int J Ophthalmol 2015;8(4):804-811.
 17. Klein BE, Klein R, Jensen SC et al. hypertension and lens opacities from the Beaver Dam eye study. Am J Ophthalmology 1995;119:640-646.
 18. Influence of Diabetes and Cardiovascular Disease on the Long-Term Incidence of Cataract: The Blue Mountains Eye Study; Jennifer S.L. Tan, Jie Jin Wang & Paul Mitchell; Ophthalmic Epidemiol. 2008 Sep-Oct; 15(5) DOI:10.1080/09286580802105806317-327.
 19. Xiaoning Yu, Danni Lyu, Xinran Dong et al; Hypertension and cataract: ametaanalysis; PLOS One; doi: 10.1371/journal.pone.0114012 December 2014.
 20. Ughade SN, et.al; Risk factors for cataract: A case control study, Indian J Ophthalmol 1998;46:221-7.
 21. Su-Ying Tsai, et.al; Epidemiologic study of age-related cataracts among an elderly Chinese population in Shih-Pai, Taiwan; Ophthalmology; June 2003;volume 110,issue 6,1089-95.
 22. Cecils Delcourt, et. al; Risk Factors for Cortical, Nuclear, and Posterior Subcapsular Cataracts The POLA Study; The POLA (Pathologies Oculaires Liées à l'Age) Am J Epidemiol. 2000 Mar 1;151(5):497-504.
 23. Nirmalan PK, Padmavathi A, Thulasiraj RD, Sex inequalities in cataract blindness burden and surgical services in South India Br J Ophthalmol 2003;87:847-9.
 24. C.O. Bekibebe, et.al. Risk factors for visually disabling age related cataracts in Ibadan; Annuals of African Medicine; Vol. 2. No.1;2003;27-32.
 25. Na K-S, Park Y-G, Han K, Mok JW, Joo C-K (2014) Prevalence of and Risk Factors for Age-Related and Anterior Polar Cataracts in a Korean Population. PLoS ONE 9(6): e96461. doi:10.1371/journal.pone.0096461.
 26. Schaumberg DA, Glynn RJ, Christensen WG; A prospective study of blood pressure and risk of cataract in men. Ann Epidemiol. 2001 Feb;11(2)104-10.
 27. M Sharma, et.al; An Epidemiological Study Of Correlates Of Cataract Among Elderly Population Aged Over 65 Years In Ut, Chandigarh; The Internet Journal of Geriatrics and Gerontology; Volume 4; Number 2.
 28. Sobti S & Sahni B; Cataract among adults aged 40 years and above in a rural area of Jammu district in India: Prevalence and Risk-factors; International J. of Healthcare & Biomedical Research, Volume: 1, Issue: 4, July 2013, Pages 284-296.
 29. Bautista LE, Vera LM, Arenas IA, Gamrara G (2005) Independent association between inflammatory markers (C-reactive protein, interleukin-6, and TNF-alpha) and essential hypertension. J Hum Hypertension 19: 149-154. doi: 10.1038/sj.jhh.1001785.
 30. Mitchell P. et.al, Medications and Cataract; The Blue Mountain Eye Study; Ophthalmology.1998 Sep;105(9):1751-8.
 31. GL Kanthan, et.al, Use of anti-hypertensive medications and topical eta-blockers and long-term incidence of cataract and cataract surgery; Br J Ophthalmol 2009;93:1210-1214, doi:10.1136/bjo.2008.153379.
 32. Klein BE, et.al, Drug use and five-year incidence of age-related cataracts: The Beaver Dam Eye Study; Ophthalmology. 2001 Sep;108(9):1670-4.
 33. Mukesh BN, et.al; Development of cataract and associated risk factors: the Visual impairment project; Arch Ophthalmol. 2006Jan;124(1):79-85.
 34. Harrison's principles of internal medicin; 18th edition;Vol2 Longo, Fanci, Kasper, Hauser, Jameson, Loscalzo;2047,2970.
 35. AK Khurana; Comprehensive ophthalmology 5th edition;599,511
 36. Systemic redox modifications in senile cataract; Virgolicci B, et.al; Rom J Intern Med 2009;47(3):279-87.
 37. Kaur J, Kukreja S, et.al; The oxidative stress in cataract patients; J Clin Diagn Res; 2012 Dec;6(10):1629-32.doi: 10.7860/JCDR/2012/4856.2626. E-pub 2012 Oct 14.