

Content available at: <https://www.ipinnovative.com/open-access-journals>

Indian Journal of Clinical and Experimental Ophthalmology

Journal homepage: www.ijceo.org

Original Research Article

Anatomical and visual outcomes after vitreoretinal surgery in paediatric retinal detachment

Chanchal Meena^{1,*}, Sanjeev K Nainiwal², Ram Swaroop Harsolia¹¹Dept. of Ophthalmology, J.L.N. Medical College and Hospital, Ajmer, Rajasthan, India²Dept. of Ophthalmology, S.M.S. Medical College and Hospital, Jaipur, Rajasthan, India

ARTICLE INFO

Article history:

Received 23-05-2023

Accepted 24-06-2023

Available online 29-09-2023

Keywords:

Paediatric vitreoretinal surgery

Retinal detachment

Pars plana vitrectomy

ABSTRACT

Background: Paediatric vitreoretinal surgery (PVRS) has unique set of challenges compared to surgery in adult eyes. An understanding of ocular growth and development is an important consideration in PVRS. Changes in ciliary body dimensions with age may affect how sclerotomies are placed, so as to avoid iatrogenic damage to the peripheral retina.

Aim: To study the visual and anatomical outcomes after vitreoretinal surgery in paediatric age group after retinal detachment.

Materials and Methods: In this prospective interventional study, 20 consecutive patients aged 18 years or younger undergoing vitreoretinal surgery for retinal detachment were included. A detailed ocular and systemic history was recorded from all patients. Pre and postoperative data were collected. All the evaluated eyes were followed up to 3 months. Outcome of surgery was measured in terms of improvement in visual acuity and retinal status.

Results: Twenty eyes of 20 patients were evaluated with male predominance (75%) and mean age of the patient was 16.4±2.01. Most common etiology was found to be trauma (65%) followed by myopia (30%). Pars plana vitrectomy (PPV) was done in 50% patients, lensectomy with vitrectomy was done in 15%, scleral buckle (SB) surgery was done in 25% cases, where as PPV-SB was done in 10% cases. Improvement in visual acuity was noted in 85% cases. Primary anatomical success was achieved in 80% cases.

Conclusion: PPV was the major surgical procedure in our study due to complexity of RD. Postoperative visual rehabilitation was also done to fight amblyopia as far as possible.

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprint@ipinnovative.com

1. Introduction

Paediatric vitreoretinal surgery (PVRS) has its unique set of challenges compared to surgery in adult eye. An understanding of ocular growth and development is an important consideration in PVRS. Changes in ciliary body dimensions with age may affect how sclerotomies are placed, so as to avoid iatrogenic damage to the peripheral retina. These changes are especially dramatic in early childhood.¹

PVRS has important differences compared to that in adults. The main indications for PVRS include rhegmatogenous retinal detachments (RRD) caused by trauma, congenital retinoschisis, marfan, stickler or morning glory syndrome, peripheral retinal degenerations, fundus coloboma, retinoblastoma, myopia, previous ocular surgery and tractional retinal detachments (TRD) caused by retinopathy of prematurity (ROP), persistent fetal vasculature (PFV), familial exudative vitreoretinopathy (FEVR) and penetrating ocular trauma. RRD in the adult usually associates with the posterior vitreous detachment (PVD), while as the RRD in paediatric age is often caused

* Corresponding author.

E-mail address: chanchalmeena986@gmail.com (C. Meena).

by trauma or congenital abnormalities. As paediatric retinal detachment (RD) usually present in late stages and paediatric ocular anatomy is featured, PVRS is more difficult and complex and has greater complication rates. The most common devastating complication of PVRS is proliferative vitreoretinopathy (PVR). In particular, creating an iatrogenic retinal break in (IRB) during PVRS can cause PVR in children because of the strong inflammatory and proliferative response.²⁻⁶

Some special evaluations such as preoperative planning, appropriate information to parents on the management, good examination and viewing of fundus under general anaesthesia, decision on application of scleral buckle (SB) or pars plana vitrectomy, the entries, angles and placement of sclerotomy and trocars, crystalline lens status, separation of the posterior hyaloid, selection of intravitreal tamponade and also visual rehabilitation are required for PVRS.⁷⁻¹⁴

2. Materials and Methods

This was a prospective interventional study of 20 consecutive patients aged 18 years or younger undergoing vitreoretinal surgery for retinal detachment at the department of Ophthalmology of Jawahar Lal Nehru Medical College, Ajmer (Rajasthan) in a time period between October 2020 to March 2022.

2.1. Inclusion criteria

1. Patients aged 18 years or younger.
2. Patients with retinal detachment who had no history of vitreoretinal surgery previously.

2.2. Exclusion criteria

1. Presence of significant corneal opacity.
2. Presence of significant conjunctival and/or scleral scarring.

A detailed ocular and systemic history was recorded from all the patients. Written informed consent was taken from all patients. Preoperative and postoperative data were collected. Descriptive statistical analysis including name, age, gender, laterality, etiology, duration of presenting symptoms, presence of ocular comorbidities, anterior segment evaluation by slit lamp biomicroscope, macular status, presence or absence of proliferative vitreoretinopathy (PVR), ocular coherence tomography (OCT)- if required, initial and final best corrected visual acuity (Snellen visual acuity converted to logMAR), type of endotamponade used, pre and postoperative lens status, intraocular pressure (IOP) measurement (by schiottz tonometer), anatomical success and complications if any. A preliminary ultrasonography was done where there was no view of fundus. Other investigations were included according to need of diagnosis and requirement of anaesthesia.

2.3. Surgical procedure

All the surgeries were performed under general anaesthesia. In patients with visible retinal breaks scleral buckle surgery was done. Three port 23G pars plana vitrectomy with silicone oil or air endotamponade was done in patients with un-detected retinal breaks, large and/or multiple retinal breaks, complicated RD with PVR and obscure ocular media. Pars plana lensectomy with vitrectomy also performed in patients with traumatic cataract. Combination of both scleral buckling and pars plana vitrectomy was also performed.

All evaluated eyes were followed up postoperatively day 1, one week, one month and three month. Outcome of surgery was measured in terms of improvement in visual acuity and retinal status.

3. Results

Demographical and clinical characteristics of patients are presented in Table 1.

Twenty eyes of 20 patients were studied. There were 15 (75%) males and 5 (25%) females. The mean age (SD) of the patients was 16.4 (2.01) years, ranging from 11 to 18 years. When coming to etiology, 13 (65%) cases were due to trauma, 6 (30%) cases due to myopia and 1 (5%) case due to fundal coloboma. Figure 1 shows the morphology of RD found in each eye. PVR was seen in 20% cases, multiple retinal holes/breaks were found in 20% cases, tractional RDs were seen in 10% cases, exudative RD in 5% cases and giant retinal tears were seen in 5% cases. Break type was not specified or known in 40% cases. When coming to status of macula; macula was off in 8 (40%) patients, whereas macula on was seen in 12 (60%) patients.

Various surgical interventions were done, and they had been chosen based on the morphology of RD. out of total 20 patients 10 (50%) patients had been treated with pars plana vitrectomy (PPV), 3 (15%) patients underwent lensectomy with vitrectomy, 5 (25%) patients were treated with scleral buckle (SB) surgery with retinopexy, and 2 (10%) patients were treated with PPV with SB. Most commonly used endotamponade was silicone oil (93.33%) and in 1 case air was used. Preoperative mean BCVA was 2.33 ± 1.12 and postoperatively at final follow up mean BCVA was 1.56 ± 0.80 with statistically significant P value (0.001) as shown in Figure 2. Visual acuity improved in 17 patients (85%), deteriorate in 2 patients (10%) and stable in one patient (5%) at 3 months. Primary anatomical success was achieved in 16 patients (80%) at final follow up.

Intraoperative complications did not occur in our study. Table 2 shows complications after surgical intervention. Early postoperative complications (≤ 1 week) included corneal edema in 2 patients, hypotony in one patient, ocular hypertension in one patient, hyphema in one patient, pre-retinal bleeding in one patient and endophthalmitis in

one patient. Late postoperative complications (>1 week) included Re-Retinal detachment (ReRD) in 3 patients and cataract in 2 patients. None of the patient developed secondary glaucoma. Corneal edema and pre-retinal bleed resolved spontaneously at one week follow up. Hyphema after surgery cleared on its own during the follow up period. Ocular hypertension was controlled by systemic and topical IOP lowering agents. Hypotony was resolved during follow up in 7 days. One patient of endophthalmitis and three patients of ReRD required second vitreoretinal surgery. Two patients who developed cataract at three months follow up required cataract extraction and IOL implantation.

Table 1: Demographic and clinical characteristics

Parameters		Number of subjects	Percentage
Gender	Male	15	75%
	Female	5	25%
Etiology	Trauma	13	65%
	Myopia	6	30%
	Fundal coloboma	1	5%
Status of macula	On	12	60%
	Off	8	40%
Endotamponade	PPV with silicone oil	14	93.33%
	PPV with air	1	6.67%




Fig. 1: Morphology of retinal detachment found in each eye

4. Discussion

This study was performed to assess the clinical characteristics of paediatric RD, surgical outcome and complications after vitreoretinal surgery in children. Out of total 20 patients, there were male predominance with 75% than females (25%). Robert Rejda et al. in his study also found higher number of male patients (75%) compared to females. This male predominance may be due to higher exposure to trauma to male gender.¹⁵




Fig. 2: Changes in mean BCVA over time

Table 2: Early and late post-operative complications

Complications	Number of subjects	Percentage
Early post-operative complications (≤1 week)	Corneal edema	2 (10%)
	Hypotony	1 (5%)
	Ocular hypertension	1 (5%)
	Hyphema	1 (5%)
	Pre-retinal bleeding	1 (5%)
	Endophthalmitis	1 (5%)
Late post-operative complications (>1 week)	Re-retinal detachment	3 (15%)
	Cataract	2 (10%)




Fig. 3: Shows scleral buckle surgery in a patient of RD



Fig. 4: Shows parsplana vitrectomy in a patient of RD




Fig. 5: B-scan image of a child showing retinal detachment with thick tractional band

Various etiological factors were present in patients with underlying retinal conditions; trauma was the most common etiology (65%) followed by myopia (30%). According to Robert Rejdak et al. trauma was the most common etiological reason (57.1%) for RD followed by high myopia (37.3%).¹⁵ Soheilian et al. in his study has evaluated clinical features, functional/anatomical outcomes after surgery in paediatric rhegmatogenous RD and found that trauma and congenital anomalies were the etiologies in paediatric rhegmatogenous RD.¹⁶ Macula was off in 40% patients. As reported by Paul C Ajamian et al. in his study that in all standard cases of macula-off RD, should get treated as quickly as possible because chronic detachments can dramatically affect the visual outcome and prognosis.¹⁷

Vitreoretinal surgeries used in the treatment of RD herein included pars plana vitrectomy, scleral buckle surgery and combination of both PPV-SB. Primary retinal reattachment rate achieved after surgery was 85% at last follow up,




Fig. 6: OCT image of a child showing retinal detachment with macula off

where as one patient of scleral buckle surgery developed endophthalmitis at one week follow up. Robert Rejdak et al. in his study achieved anatomical success in 86% cases.¹⁵ Gurler et al. also retinal reattachment after primary surgery in about 80% of cases. He reported better BCVA with RD associated with trauma than in cases of myopia.¹⁸

The surgical principles that guide repair of adult retinal detachment also direct the repair of paediatric retinal detachment, relief of vitreous traction on the causative retinal defects, reapproximation of the retina to the retinal pigment epithelium.¹⁹ Paediatric vitreous is often more rigid and adherent than adult vitreous. In cases with broad peripheral pathology, the vitreous itself can help to tamponade a break, once the traction has been relieved with a scleral buckle, allowing resorption of sub-retinal fluid.¹² Vitreous that is retained near a break may contract, leading to recurrent detachment. Higher intraocular cellular activity means that the probability of developing PVR and recurrent detachment is higher in paediatric patients with retinal detachment when compared to adults with the same.¹⁹ Vitreoretinal surgeries are challenging in children due to strong vitreoretinal adhesions so that iatrogenic retinal break easily occur. It can be associated with cataract formation due to lens injury during surgery. Subsequent cataract surgery was required in 2 cases as described above.

In this study mean BCVA was improved in 85% cases, deteriorate in 10%, and stable in 5% cases. Good visual outcome after surgery was favourably influenced by preoperative visual acuity. The paediatric traumatic RD usually present early while no traumatic RD usually present at a later stage compared to adults. Therefore the management often lasts for a long because to fight the amblyopia and visual rehabilitation compared to adults.

5. Conclusion

In this study, mean age of the patients was 16.4±2.01 with male predominance. Most common etiology was found to be trauma followed by myopia. PPV was the major surgical procedure in this study due to complexity of RD. Most common endotamponade used was silicone oil in PPV. The visual and anatomical outcome following surgery were favourable in 85% and 80% cases respectively. Four eye underwent second vitreoretinal surgery (3 patients of ReRD and 1 patient of endophthalmitis) and two eyes for cataract surgery. Postoperative visual rehabilitation was also done to fight to amblyopia as far as possible.

6. Source of Funding

None.

7. Conflict of Interest

None.

References

- Lemley CA, Han DP. An age-based method for planning sclerotomy placement during pediatric vitrectomy: a 12-year experience. *Trans Am Ophthalmol Soc.* 2007;105:86–9.
- Al-Zaaidi S, Al-Rashaed S, AL-Harthi E, AL-Kahtani E, El-Asrar AA. Rhegmatogenous retinal detachment in children 16 years of age or younger. *Clin Ophthalmol.* 2013;7:1001–14.
- Wadhwa N, Venkatesh P, Sampangi R, Garg S. Rhegmatogenous retinal detachments in children in India: clinical characteristics, risk factors, and surgical outcomes. *J AAPOS.* 2008;12(6):551–4.
- Wenick AS, Barañano DE. Evaluation and management of pediatric rhegmatogenous retinal detachment. *Saudi J Ophthalmol.* 2012;26(3):255–63.
- Read SP, Aziz HA, Kuriyan A. Retinal detachment surgery in a pediatric population: visual and anatomic outcomes. *Retina.* 2018;38(7):1393–1402.
- Gan NY, Lam WC. Retinal detachments in the pediatric population. *Taiwan J Ophthalmol.* 2018;8(4):222–6.
- Yonekawa Y, Fine HF. Practical Pearls in Pediatric Vitreoretinal Surgery. *Ophthalmic Surg Lasers Imaging Retina.* 2018;49(8):561–5.
- Leung EH, Berrocal AM. Pediatric Microincision Vitreoretinal Surgery. *Int Ophthalmol Clin.* 2016;56(4):203–8.
- Hu Y, Tang S. Major challenges in vitreoretinal surgery. *Taiwan J Ophthalmol.* 2014;5(1):9–14.
- Sebag J. Age-related differences in the human vitreoretinal interface. *Arch Ophthalmol.* 1991;109(7):966–971.
- Ranchod TM, Capone A. Tips and tricks in pediatric vitreoretinal surgery. *Int Ophthalmol Clin.* 2011;51(1):173–83.
- Baumal CR, Berrocal AM. Pearls for Pediatric Retinal Surgery; 2018. Available from: <https://retinatoday.com/articles/2018-july-aug/pearls-for-pediatric-retinal-surgery>.
- Nudleman E, Capone A. Management of Complicated Vitreoretinal Cases in Children. In: Patelli F, Stanislao R, editors. Management of Complicated Vitreoretinal Diseases. Switzerland: Springer International Publishing; 2015. p. 21–30.
- Wright LM, Harper CA, Chang EY. Management of infantile and Childhood Retinopathies: Optimized Pediatric Pars Plana Vitrectomy Sclerotomy Nomogram. *Ophthalmol Retina.* 2018;2(12):1227–34.
- Rejda R, Nowakowska D, Wrona K, Maciejewski R, Junemann AG, Nowomiejska K. Outcomes of Vitrectomy in Pediatric Retinal Detachment with Proliferative Vitreoretinopathy. *J Ophthalmol.* 2017;2017:8109390. doi:10.1155/2017/8109390.
- Soheilian M, Ramezani A, Malihi M, Yaseri M, Ahmadi H, Dehghan MH, et al. Clinical features and surgical outcomes of pediatric rhegmatogenous retinal detachment. *Retina.* 2009;29(4):545–51.
- Ajamian PC. Off to a bad start: The timing of surgical intervention for macula-off RD is debatable, and COVID could complicate things further; 2020. Available from: <https://www.reviewofoptometry.com/article/off-to-a-bad-start>.
- Gurler B, Coskun E, Öner V, Comez A, Erbagci I. Clinical characteristics and surgical outcomes of pediatric rhegmatogenous retinal detachment. *Int Ophthalmol.* 2016;36(4):521–5.
- Holladay JT. Proper method for calculating average visual acuity. *J Refract Surg.* 1997;13(4):388–91.

Author biography

Chanchal Meena, Junior Resident  <https://orcid.org/0009-0002-4774-3891>

Sanjeev K Nainiwal, Senior Professor and Unit Head

Ram Swaroop Harsolia, Associate Professor

Cite this article: Meena C, Nainiwal SK, Harsolia RS. Anatomical and visual outcomes after vitreoretinal surgery in paediatric retinal detachment. *Indian J Clin Exp Ophthalmol* 2023;9(3):403–407.